

CLASSICAL YANG-BAXTER EQUATION AND LOW DIMENSIONAL TRIANGULAR LIE BIALGEBRAS *

Shouchuan Zhang

Department of Mathematics, Nanjing University, 210008
P.R.China

Abstract

All solutions of constant classical Yang-Baxter equation (CYBE) in Lie algebra L with $\dim L \leq 3$ are obtained and the sufficient and necessary conditions which $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary (or triangular) Lie bialgebra are given. The strongly symmetric elements in $L \otimes L$ are found and they all are solutions of CYBE in L with $\dim L \leq 3$.

0 Introduction

The concept and structures of Lie coalgebras were introduced and studied by W.Michaelis in [6] [7]. V.G.Drinfel'd and A.A.Belavin in [2] [1] introduced the notion of triangular, coboundary Lie bialgebra L associated to a solution $r \in L \otimes L$ of the CYBE and gave a classification of solutions of CYBE with parameter for simple Lie algebras. W.Michaelis in [5] obtained the structure of a triangular, coboundary Lie bialgebra on any Lie algebra containing linearly independent elements a and b satisfying $[a, b] = \alpha b$ for some non-zero $\alpha \in k$ by setting $r = a \otimes b - b \otimes a$.

The Yang-Baxter equation first came up in a paper by Yang as factorization condition of the scattering S-matrix in the many-body problem in one dimension and in work of Baxter

*This work is supported by National Science Foundation

on exactly solvable models in statistical mechanics. It has been playing an important role in mathematics and physics (see [1] , [9]). Attempts to find solutions of The Yang-Baxter equation in a systematic way have let to the theory of quantum groups. The Yang-Baxter equation is of many forms. The classical Yang-Baxter equation is one.

In many applications one need to know the solutions of classical Yang-Baxter equation and know if a Lie algebra is a coboundary Lie bialgebra or a triangular Lie bialgebra. A systematic study of low dimensional Lie algebras, specially, of those Lie algebras that play a role in physics (as e.g. $sl(2, \mathbf{C})$, or the Heisenberg algebra), is very useful.

In this paper, we obtain all solutions of constant classical Yang-Baxter equation (CYBE) in Lie algebra L with $\dim L \leq 3$ and give the sufficient and necessary conditions which $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary (or triangular) Lie bialgebra. We find the strongly symmetric elements in $L \otimes L$ and show they all are solutions of CYBE in L with $\dim L \leq 3$. Using these conclusions, we study the Lie algebra $sl(2)$.

In order to make the paper somewhat self-contained, we begin by recalling the definition of Lie bialgebra (all of the following definitions in the introduction are the same as [5, P368-371]).

Let k be a field with $\text{char } k \neq 2$ and L be a Lie algebra over k . If $r = \sum a_i \otimes b_i \in L \otimes L$ and $x \in L$, we define

$$\begin{aligned} [r^{12}, r^{13}] &:= \sum_{i,j} [a_i, a_j] \otimes b_i \otimes b_j \\ [r^{12}, r^{23}] &:= \sum_{i,j} a_i \otimes [b_i, a_j] \otimes b_j \\ [r^{13}, r^{23}] &:= \sum_{i,j} a_i \otimes a_j \otimes [b_i, b_j] \\ x \cdot r &:= \sum_i [x, a_i] \otimes b_i + a_i \otimes [x, b_i] \\ \Delta_r(x) &:= x \cdot r \end{aligned}$$

and call

$$[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}] = 0$$

the classical Yang-Baxter equation (CYBE).

Let

$$\tau : L \otimes L \longrightarrow L \otimes L$$

denote the natural twist map (defined by $x \otimes y \mapsto y \otimes x$), and let

$$\xi : L \otimes L \otimes L \longrightarrow L \otimes L \otimes L$$

the map defined by $x \otimes y \otimes z \mapsto y \otimes z \otimes x$ for any $x, y, z \in L$.

a vector space L is called a Lie coalgebra, if there exists a linear map

$$\Delta : L \longrightarrow L \otimes L$$

such that

- (i) $Im\Delta \subseteq Im(1 - \tau)$ and
 - (ii) $(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta = 0$
- a vector space $(L, [\], \Delta)$ is called a Lie bialgebra if
- (i) $(L, [\])$ is a Lie algebra,
 - (ii) (L, Δ) is a Lie coalgebra,
 - (iii) for all $x, y \in L$

$$\Delta[x, y] = x \cdot \Delta(y) - y \cdot \Delta(x),$$

where, for all $x, a_i, b_i \in L$,

$(L, [\], \Delta, r)$ is called a coboundary Lie bialgebra, if $(L, [\], \Delta)$ is a Lie bialgebra and $r \in Im(1 - \tau) \subseteq L \otimes L$ such that

$$\Delta(x) = x \cdot r$$

for all $x \in L$. a coboundary Lie bialgebra $(L, [\], \Delta, r)$ is called triangular, if r is a solution of CYBE.

1 The solutions of CYBE

In this section, we find the general solution of CYBE for Lie algebra L with $dim L \leq 3$.

Jacobson gave a classification of Lie algebras with their dimension $dim L \leq 3$ in [3, P 11-14]. We now write all of their operation as follows:

- (I) If L is an abelian Lie algebra, then its operation is trivial;
- (II) If $dim L = dim L' = 3$, then there exist a basis $\{e_1, e_2, e_3\}$ and $\alpha, \beta \in k$ with $\alpha\beta \neq 0$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$;
- (III) If $dim L = 3$ and $L' \subseteq$ the center of L with $dim L' = 1$, then there exist a basis $\{e_1, e_2, e_3\}$ and $\alpha, \beta \in k$ with $\alpha = \beta = 0$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$;
- (IV) If k is an algebraically closed field and $dim L = 3$ with $dim L' = 2$, then there exist a basis $\{e_1, e_2, e_3\}$ and $\beta, \delta \in k$ with $\delta \neq 0$ such that $[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2$;
- (V) If $dim L = 3$ and $L' \not\subseteq$ the center of L with $dim L' = 1$, then there exist a basis $\{e_1, e_2, e_3\}$ and $\beta, \delta \in k$ with $\beta = \delta = 0$ such that $[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2$;
- (VI) If $dim L = 2$ with $dim L' = 1$, then there exists a basis $\{e, f\}$ such that $[e, f] = e$.

We like to point out the actual meaning of the different algebras, namely (II) includes $sl(2, \mathbf{C})$ and its real form $su(2)$, (III) is just the three-dimensional Heisenberg algebra, (IV) includes the (1+1) Poincare algebra, (V) is the (one-dimensional) central extension of the non-abelian two-dimensional Lie algebra and (VI) is the non-abelian two-dimensional Lie algebra itself.

For case (I), the two treated problems are trivial.

Definition 1.1 Let $\{e_1, e_2, \dots, e_n\}$ be a basis of vector space V and $r = \sum_{i,j=1}^n k_{ij}(e_i \otimes e_j) \in V \otimes V$, where $k_{ij} \in k$, for $i, j = 1, 2, \dots, n$.

(i) If

$$k_{ij} = k_{ji} \quad k_{ij}k_{lm} = k_{il}k_{jm}$$

for $i, j, l, m = 1, 2, \dots, n$, then r is called strongly symmetric to the basis $\{e_1, e_2, \dots, e_n\}$;

(ii) If $k_{ij} = -k_{ji}$, for $i, j = 1, 2, \dots, n$, then r is called skew symmetric.

(iii) Let $\dim V = 3$ and $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v, \alpha, \beta \in k$. If $p = -q, s = -t, u = -v, x = \alpha z, y = \beta z$ and

$$\alpha\beta z^2 + \beta s^2 + \alpha u^2 + p^2 = 0,$$

then r is called α, β -skew symmetric to the basis $\{e_1, e_2, e_3\}$.

Obviously, skew symmetry does not depend on the particular choice of basis of V . We need to know if strong symmetry depend on the particular choice of basis of V .

Lemma 1.2 (I) If V is a finite-dimensional vector space, then the strong symmetry does not depend on the particular choice of basis of V ;

(II) Let $\{e_1, e_2, e_3\}$ be a basis of vector space V , $p, q, s, t, u, v, x, y, z \in k$ and $r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$.

Then r is strongly symmetric iff

$$xy = p^2, xz = s^2, yz = u^2, xu = sp, ys = pu, zp = su, p = q, s = t, u = v$$

iff

$$xy = p^2, xz = s^2, yz = u^2, xu = sp, p = q, s = t, u = v$$

iff

$$xy = p^2, xz = s^2, yz = u^2, ys = up, p = q, s = t, u = v$$

iff

$$xy = p^2, xz = s^2, yz = u^2, zp = su, p = q, s = t, u = v$$

iff

$$\begin{aligned} r &= z^{-1}su(e_1 \otimes e_2) + z^{-1}su(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + u(e_3 \otimes e_2) + \\ &z^{-1}s^2(e_1 \otimes e_1) + z^{-1}u^2(e_2 \otimes e_2) + z(e_3 \otimes e_3) \text{ with } z \neq 0; \text{ or,} \\ r &= p(e_1 \otimes e_2) + p(e_2 \otimes e_1) + x(e_1 \otimes e_1) + x^{-1}p^2(e_2 \otimes e_2) \text{ with } x \neq 0; \text{ or,} \\ r &= y(e_2 \otimes e_2) \end{aligned}$$

Proof. (I) Let $\{e_1, e_2, \dots, e_n\}$ and $\{e'_1, e'_2, \dots, e'_n\}$ are two basis of V and $r = \sum_{i,j=1}^n k_{ij}(e_i \otimes e_j)$ be strongly symmetric to the basis $\{e_1, e_2, \dots, e_n\}$. It is sufficient to show that r is strongly symmetric to the basis $\{e'_1, e'_2, \dots, e'_n\}$. Obviously, there exists $q_{ij} \in k$ such that $e_i = \sum_s e'_s q_{si}$ for $i = 1, 2, \dots, n$. By computation, we have that

$$r = \sum_{s,t} \left(\sum_{i,j} k_{ij} q_{si} q_{tj} \right) (e_s \otimes e_t).$$

If set $k'_{st} = \sum_{i,j} k_{ij} q_{si} q_{tj}$, then for $l, m, u, v = 1, 2, \dots, n$, we have that

$$\begin{aligned} k'_{lm} k'_{uv} &= \sum_{i,j,s,t} k_{ij} k_{st} q_{li} q_{mj} q_{us} q_{vt} \\ &= \sum_{i,j,s,t} k_{is} k_{jt} q_{li} q_{mj} q_{us} q_{vt} \quad (\text{by } k_{ij} k_{st} = k_{is} k_{jt}) \\ &= k'_{lu} k'_{mv} \end{aligned}$$

and

$$k'_{lm} = k'_{ml}$$

which implies that r is strongly symmetric to the basis $\{e'_1, e'_2, \dots, e'_n\}$

(II) We only show that if

$$xy = p^2, xz = s^2, yz = u^2, xu = sp, p = q, s = t, u = v,$$

then

$$ys = pu, zp = su.$$

If $p \neq 0$, then $ysp = yxu = up^2$ and $usp = uxu = xyz = zp^2$, which implies $ys = up$ and $us = zp$. If $p = 0$, then $x = 0$ or $y = 0$, which implies $s = 0$ or $u = 0$. Consequently, $ys = up$ and $us = zp$. $\square$

Proposition 1.3 *Let L be a Lie algebra with $\dim L = 2$. Then r is a solution of CYBE iff r is strongly symmetric or skew symmetric.*

Proof. It is not hard since the tensor r has only 4 coefficients. $\square$

Proposition 1.4 *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$, where $\alpha, \beta \in k$. Let $p, q, s, t, u, v, x, y, z \in k$.*

(I) If r is strongly symmetric or α, β -skew symmetric to basis $\{e_1, e_2, e_3\}$, then r is a solution of CYBE;

(II) If $\alpha \neq 0, \beta \neq 0$, then r is a solution of CYBE in L iff r is strongly symmetric or α, β -skew symmetric;

(III) If $\alpha = \beta = 0$, then r is a solution of CYBE in L iff

$$r = p(e_1 \otimes e_2) + p(e_2 \otimes e_1) + s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

with $p \neq 0, p^2 = xy, xu = sp, xv = tp, tu = vs$, or

$$r = s(e_1 \otimes e_3) + t(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

with $xy = xu = xv = ys = yt = 0$ and $tu = vs$.

Proof . Let $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) \in L \otimes L$ and $k_{ij} \in k$, with $i, j = 1, 2, 3$. It is clear that

$$\begin{aligned} [r^{12}, r^{13}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} [e_i, e_s] \otimes e_j \otimes e_t \\ [r^{12}, r^{23}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} e_i \otimes [e_j, e_s] \otimes e_t \\ [r^{13}, r^{23}] &= \sum_{i,j=1}^3 \sum_{s,t=1}^3 k_{ij} k_{st} e_i \otimes e_s \otimes [e_j, e_t] \end{aligned}$$

By computation, for all $i, j, n = 1, 2, 3$, we have that the coefficient of $e_j \otimes e_i \otimes e_i$ in $[r^{12}, r^{13}]$ is zero and $e_i \otimes e_i \otimes e_j$ in $[r^{13}, r^{23}]$ is zero.

We now see the coefficient of $e_i \otimes e_j \otimes e_n$

in $[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}]$.

- (1) $e_1 \otimes e_1 \otimes e_1(\alpha k_{12} k_{31} - \alpha k_{13} k_{21})$
- (2) $e_2 \otimes e_2 \otimes e_2(-\beta k_{21} k_{32} + \beta k_{23} k_{12})$
- (3) $e_3 \otimes e_3 \otimes e_3(k_{31} k_{23} - k_{32} k_{13})$
- (4) $e_1 \otimes e_2 \otimes e_3(\alpha k_{22} k_{33} - \alpha k_{32} k_{23} - \beta k_{11} k_{33} + \beta k_{13} k_{13} + k_{11} k_{22} - k_{12} k_{21})$
- (5) $e_2 \otimes e_3 \otimes e_1(\beta k_{33} k_{11} - \beta k_{13} k_{31} - k_{22} k_{11} + k_{21} k_{21} + \alpha k_{22} k_{33} - \alpha k_{23} k_{32})$
- (6) $e_3 \otimes e_1 \otimes e_2(k_{11} k_{22} - k_{21} k_{12} - \alpha k_{33} k_{22} + \alpha k_{32} k_{32} + \beta k_{33} k_{11} - \beta k_{31} k_{13})$
- (7) $e_1 \otimes e_3 \otimes e_2(-\alpha k_{33} k_{22} + \alpha k_{23} k_{32} + k_{11} k_{22} - k_{12} k_{12} - \beta k_{11} k_{33} + \beta k_{13} k_{31})$
- (8) $e_3 \otimes e_2 \otimes e_1(-k_{22} k_{11} + k_{12} k_{21} + \beta k_{33} k_{11} - \beta k_{31} k_{31} - \alpha k_{33} k_{22} + \alpha k_{32} k_{23})$
- (9) $e_2 \otimes e_1 \otimes e_3(-\beta k_{11} k_{33} + \beta k_{31} k_{13} + \alpha k_{22} k_{33} - \alpha k_{23} k_{23} - k_{22} k_{11} + k_{21} k_{12})$
- (10) $e_1 \otimes e_1 \otimes e_2(-\alpha k_{31} k_{22} + \alpha k_{21} k_{32} + \alpha k_{12} k_{32} - \alpha k_{13} k_{22})$

- (11) $e_2 \otimes e_1 \otimes e_1(-\alpha k_{23}k_{21} + \alpha k_{22}k_{31} + \alpha k_{22}k_{13} - \alpha k_{23}k_{12})$
- (12) $e_1 \otimes e_1 \otimes e_3(\alpha k_{21}k_{33} - \alpha k_{31}k_{23} + \alpha k_{12}k_{33} - \alpha k_{13}k_{23})$
- (13) $e_3 \otimes e_1 \otimes e_1(\alpha k_{32}k_{31} - \alpha k_{33}k_{21} + \alpha k_{32}k_{13} - \alpha k_{33}k_{12})$
- (14) $e_2 \otimes e_2 \otimes e_1(-\beta k_{12}k_{31} + \beta k_{32}k_{11} + \beta k_{23}k_{11} - \beta k_{21}k_{31})$
- (15) $e_1 \otimes e_2 \otimes e_2(-\beta k_{11}k_{32} + \beta k_{13}k_{12} - \beta k_{11}k_{23} + \beta k_{13}k_{21})$
- (16) $e_2 \otimes e_2 \otimes e_3(\beta k_{32}k_{13} - \beta k_{12}k_{33} - \beta k_{21}k_{33} + \beta k_{23}k_{13})$
- (17) $e_3 \otimes e_2 \otimes e_2(-\beta k_{31}k_{32} + \beta k_{33}k_{12} + \beta k_{33}k_{21} - \beta k_{31}k_{23})$
- (18) $e_3 \otimes e_3 \otimes e_1(k_{13}k_{21} - k_{23}k_{11} + k_{31}k_{21} - k_{32}k_{11})$
- (19) $e_3 \otimes e_3 \otimes e_2(-k_{23}k_{12} + k_{13}k_{22} - k_{32}k_{12} + k_{31}k_{22})$
- (20) $e_2 \otimes e_3 \otimes e_3(k_{21}k_{23} - k_{22}k_{13} + k_{21}k_{32} - k_{22}k_{31})$
- (21) $e_1 \otimes e_3 \otimes e_3(-k_{12}k_{13} + k_{11}k_{23} - k_{12}k_{31} + k_{11}k_{32})$
- (22) $e_1 \otimes e_3 \otimes e_1(\alpha k_{23}k_{31} - \alpha k_{33}k_{21} - k_{12}k_{11} + k_{11}k_{21} + \alpha k_{12}k_{33} - \alpha k_{13}k_{32})$
- (23) $e_1 \otimes e_2 \otimes e_1(-\alpha k_{32}k_{21} + \alpha k_{22}k_{31} - \beta k_{11}k_{31} + \beta k_{13}k_{11} - \alpha k_{13}k_{22} + \alpha k_{12}k_{23})$
- (24) $e_2 \otimes e_1 \otimes e_2(\beta k_{31}k_{12} - \beta k_{11}k_{32} + \alpha k_{22}k_{32} - \alpha k_{23}k_{22} - \beta k_{21}k_{13} + \beta k_{23}k_{11})$
- (25) $e_2 \otimes e_3 \otimes e_2(-\beta k_{13}k_{32} + \beta k_{33}k_{12} + k_{21}k_{22} - k_{22}k_{12} - \beta k_{21}k_{33} + \beta k_{23}k_{31})$
- (26) $e_3 \otimes e_2 \otimes e_3(k_{12}k_{23} - k_{22}k_{13} - \beta k_{31}k_{33} + \beta k_{33}k_{13} + k_{31}k_{22} - k_{32}k_{21})$
- (27) $e_3 \otimes e_1 \otimes e_3(-k_{21}k_{13} + k_{11}k_{23} + \alpha k_{32}k_{33} - \alpha k_{33}k_{23} + k_{31}k_{12} - k_{32}k_{11}).$

Let $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v$.

It follows from (1)-(27) that

- (28) $\alpha pt = \alpha qs$
- (29) $\beta qv = \beta pu$
- (30) $tu = vs$
- (31) $\alpha yz - \beta xz + xy - \alpha uv + \beta s^2 - pq = 0.$
- (32) $\beta zx - yx + \alpha yz - \beta st + q^2 - \alpha uv = 0$
- (33) $xy - \alpha zy + \beta zx - pq + \alpha v^2 - \beta st = 0$
- (34) $-\alpha zy + xy - \beta xz + \alpha uv - p^2 + \beta st = 0$
- (35) $-xy + \beta xz - \alpha yz + pq - \beta t^2 + \alpha uv = 0$
- (36) $-\beta xz + \alpha yz - yx + \beta st - \alpha u^2 + pq = 0$
- (37) $\alpha(-ty + qv + pv - sy) = 0$
- (38) $\alpha(-uq + yt + ys - up) = 0$
- (39) $\alpha(qz - tu + pz - su) = 0$
- (40) $\alpha(vt - zq + vs - zp) = 0$
- (41) $\beta(-pt + vx + ux - qt) = 0$
- (42) $\beta(-xv + sp - xu + sq) = 0$
- (43) $\beta(vs - pz + us - qz) = 0$
- (44) $\beta(-tv + zp + zq - tu) = 0$

$$\begin{aligned}
(45) \quad & sq - ux + tq - vx = 0 \\
(46) \quad & -up + sy - vp + ty = 0 \\
(47) \quad & qu - ys + qv - yt = 0 \\
(48) \quad & -ps + xu - pt + xv = 0. \\
(49) \quad & \alpha ut - \alpha zq - px + xq + \alpha pz - \alpha sv = 0 \\
(50) \quad & -\alpha vq + \alpha yt - \beta xt + \beta sx - \alpha sy + \alpha pu = 0 \\
(51) \quad & \beta tp - \beta xv + \alpha yv - \alpha uy - \beta qs + \beta ux = 0 \\
(52) \quad & -\beta sv + \beta zp + qy - yp - \beta qz + \beta ut = 0 \\
(53) \quad & pu - ys - \beta tz + \beta zs + ty - vq = 0 \\
(54) \quad & -qs + xu + \alpha vz - \alpha zu + tp - vx = 0
\end{aligned}$$

It is clear that r is the solution of CYBE iff relations (28)-(54) hold.

(I) By computation, we have that if r is strongly symmetric or α, β -skew symmetric then relations (28)-(54) and so r is a solution of CYBE;

(II) Let $\alpha \neq 0$ and $\beta \neq 0$. By part (I), we only need show that if r is a solution of CYBE, then r is strongly symmetric or α, β -skew symmetric.

By computation, we have

$$(55) \quad q^2 = p^2 \text{ (by (34) + (32))};$$

$$(56) \quad u^2 = v^2 \text{ (by (33) + (36))};$$

$$(57) \quad t^2 = s^2 \text{ (by (31) + (35))};$$

(a). If $s = t \neq 0$, then we have that

$$p = q \text{ by (28)}$$

$$u = v \text{ by (30)}$$

$$(58) \quad yz = u^2 \text{ by (35) - (32)}$$

$$(59) \quad xz = s^2 \text{ by (32) + (33)}$$

$$(60) \quad xy = p^2 \text{ by (36) - (31)}$$

$$(61) \quad xu = sp \text{ by (41)}$$

then r is strongly symmetric.

Similarly, we can show that if $p = q \neq 0$, or $u = v \neq 0$, then r is strongly symmetric.

(b). If $s = -t \neq 0$, then we have that

$$p = -q \text{ by (28)}$$

$$u = -v \text{ by (30)}$$

$$y = \beta z \text{ by (53)}$$

$$x = \alpha z \text{ by (50)}$$

$$\alpha\beta z^2 + \alpha u^2 + \beta s^2 + p^2 = 0 \text{ by (31)}$$

Then r is α, β -skew symmetric.

Similarly, we have that if $p = -q \neq 0$ or $u = -v \neq 0$, then r is α, β -skew symmetric.

(c). If $s = t = u = v = p = q = 0$, then we have that

$$xz = 0 \text{ by (32) + (33)}$$

$$yz = 0 \text{ by (32) + (31)}$$

$$xy = 0 \text{ by (31)}$$

Thus r is strongly symmetric.

(III) Let $\alpha = \beta = 0$. It is clear that system of equations (28)-(54) is equivalent to the below

$$\begin{cases} tu = vs, q = p, p^2 = xy \\ (s+t)p - (u+v)x = 0 \\ (s+t)y - (u+v)p = 0 \\ (t-s)y + (u-v)p = 0 \\ (t-s)p + (u-v)x = 0 \end{cases}$$

It is easy to check that if r is one case in part (III) then the system of equations hold. Conversely, if r is a solution of CYBE, we shall show that r is one of two cases in part (III). If $p \neq 0$ then r is the first case in part (III). If $p = 0$ then r is the second case in part (III). $\square$

In particular, Proposition 1.4 implies:

Example 1.5 *Let*

$$sl(2) := \{x \mid x \text{ is a } 2 \times 2 \text{ matrix with trace zero over } k\}$$

and

$$e_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, e_3 = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix}.$$

Thus L is a Lie algebra (defined by $[x, y] = xy - yx$) with a basis $\{e_1, e_2, e_3\}$. It is clear that

$$[e_1, e_2] = e_3, [e_2, e_3] = 4e_1, [e_3, e_1] = -4e_2.$$

Consequently, r is a solution of CYBE iff r is strongly symmetric or 4, -4-skew symmetric to the basis $\{e_1, e_2, e_3\}$.

Proposition 1.6 *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2$, where $\beta, \delta \in k$. Let $p, q, s, t, u, v, x, y, z \in k$.*

(I) If r is strongly symmetric, then r is a solution of CYBE.

(II) If $\beta = 0, \delta \neq 0$, then r is a solution of CYBE in L iff r is strongly symmetric, or

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$$

with $xu = xs = ys = yu = (1 - \delta)us = (1 + \delta)s(q + p) = (1 + \delta)u(q + p) = 0$

(III) If $\beta \neq 0$ and $\delta = 1$, then r is a solution of CYBE in L iff r is strongly symmetric, or

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$$

with $xu = yu = u(q + p) = 0$

(IV) If $\beta = \delta = 0$, then r is a solution of CYBE in L iff

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) + s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2) + z(e_3 \otimes e_3)$$

with $z \neq 0, zp = vs, zq = us, zx = s^2$, or

$$r = p(e_1 \otimes e_2) + q(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) + v(e_3 \otimes e_2) + x(e_1 \otimes e_1) + y(e_2 \otimes e_2)$$

with $us = vs = xs = xu = xv = 0, up = qv, s(p + q) = 0$

Proof. Let $r = \sum_{i,j=1}^3 k_{ij}(e_i \otimes e_j) \in L \otimes L$ and $k_{ij} \in k$, with $i, j = 1, 2, 3$. By computation, for all $i, j, n = 1, 2, 3$, we have that the coefficient of $e_j \otimes e_i \otimes e_i$ in $[r^{12}, r^{13}]$ is zero and $e_i \otimes e_i \otimes e_j$ in $[r^{13}, r^{23}]$ is zero.

We now see the coefficient of $e_i \otimes e_j \otimes e_n$

in $[r^{12}, r^{13}] + [r^{12}, r^{23}] + [r^{13}, r^{23}]$.

- (1) $e_1 \otimes e_1 \otimes e_1(-k_{13}k_{11} + k_{11}k_{31})$;
- (2) $e_2 \otimes e_2 \otimes e_2(-\beta k_{23}k_{12} + \beta k_{21}k_{32} - \delta k_{23}k_{22} + \delta k_{22}k_{32})$;
- (3) $e_3 \otimes e_3 \otimes e_3(0)$;
- (4) $e_1 \otimes e_2 \otimes e_3(-k_{32}k_{13} + k_{12}k_{33} - \beta k_{13}k_{13} + \beta k_{11}k_{33} - \delta k_{13}k_{23} + \delta k_{12}k_{33})$;
- (5) $e_2 \otimes e_3 \otimes e_1(-\beta k_{33}k_{11} + \beta k_{13}k_{31} - \delta k_{33}k_{21} + \delta k_{23}k_{31} - k_{23}k_{31} + k_{21}k_{33})$;
- (6) $e_3 \otimes e_1 \otimes e_2(-k_{33}k_{12} + k_{31}k_{32} - \beta k_{33}k_{11} + \beta k_{31}k_{13} - \delta k_{33}k_{12} + \delta k_{32}k_{13})$;
- (7) $e_1 \otimes e_3 \otimes e_2(-k_{33}k_{12} + k_{13}k_{32} - \beta k_{13}k_{32} + \beta k_{11}k_{33} - \delta k_{13}k_{32} + \delta k_{12}k_{33})$;
- (8) $e_3 \otimes e_2 \otimes e_1(-\beta k_{33}k_{11} + \beta k_{31}k_{31} - \delta k_{33}k_{21} + \delta k_{32}k_{31} - k_{33}k_{21} + k_{31}k_{23})$;
- (9) $e_2 \otimes e_1 \otimes e_3(-\beta k_{31}k_{13} + \beta k_{11}k_{33} - \delta k_{31}k_{23} + \delta k_{21}k_{33} - k_{23}k_{13} + k_{21}k_{33})$;
- (10) $e_1 \otimes e_1 \otimes e_2(-k_{31}k_{12} + k_{11}k_{32} - k_{13}k_{12} + k_{11}k_{32})$;
- (11) $e_2 \otimes e_1 \otimes e_1(-k_{23}k_{11} + k_{21}k_{31} - k_{23}k_{11} + k_{21}k_{13})$;
- (12) $e_1 \otimes e_1 \otimes e_3(-k_{31}k_{13} + k_{11}k_{33} - k_{13}k_{13} + k_{11}k_{33})$;

- (13) $e_3 \otimes e_1 \otimes e_1(-k_{33}k_{11} + k_{31}k_{31} - k_{33}k_{11} + k_{31}k_{13});$
- (14) $e_2 \otimes e_2 \otimes e_1(-\beta k_{32}k_{11} + \beta k_{12}k_{31} - \delta k_{32}k_{21} + \delta k_{22}k_{31} - \beta k_{23}k_{11} + \beta k_{21}k_{31} - \delta k_{23}k_{21} + \delta k_{22}k_{31});$
- (15) $e_1 \otimes e_2 \otimes e_2(-\beta k_{13}k_{12} + \beta k_{11}k_{32} - \delta k_{13}k_{22} + \delta k_{12}k_{32} - \beta k_{13}k_{21} + \beta k_{11}k_{23} - \delta k_{13}k_{22} + \delta k_{12}k_{23});$
- (16) $e_2 \otimes e_2 \otimes e_3(-\beta k_{32}k_{13} + \beta k_{12}k_{33} - \delta k_{32}k_{23} + \delta k_{22}k_{33} - \beta k_{23}k_{13} + \beta k_{21}k_{33} - \delta k_{23}k_{23} + \delta k_{22}k_{33});$
- (17) $e_3 \otimes e_2 \otimes e_2(-\beta k_{33}k_{12} + \beta k_{31}k_{32} - \delta k_{33}k_{22} + \delta k_{32}k_{32} - \beta k_{33}k_{21} + \beta k_{31}k_{23} - \delta k_{33}k_{22} + \delta k_{32}k_{23});$
- (18) $e_3 \otimes e_3 \otimes e_1(0);$
- (19) $e_1 \otimes e_3 \otimes e_3(0);$
- (20) $e_3 \otimes e_3 \otimes e_2(0);$
- (21) $e_2 \otimes e_3 \otimes e_3(0);$
- (22) $e_1 \otimes e_3 \otimes e_1(-k_{33}k_{11} + k_{13}k_{31} - k_{13}k_{31} + k_{11}k_{33}) = e_1 \otimes e_3 \otimes e_1(0);$
- (23) $e_1 \otimes e_2 \otimes e_1(-k_{32}k_{11} + k_{12}k_{31} - \beta k_{13}k_{11} + \beta k_{11}k_{31} - \delta k_{13}k_{21} + \delta k_{12}k_{31} - k_{13}k_{21} + k_{11}k_{23});$
- (24) $e_2 \otimes e_1 \otimes e_2(-\beta k_{31}k_{12} + \beta k_{11}k_{32} - \delta k_{31}k_{22} + \delta k_{21}k_{32} - k_{23}k_{12} + k_{21}k_{32} - \beta k_{23}k_{11} + \beta k_{21}k_{13} - \delta k_{23}k_{12} + \delta k_{22}k_{13});$
- (25) $e_2 \otimes e_3 \otimes e_2(-\beta k_{33}k_{12} + \beta k_{13}k_{32} - \delta k_{33}k_{22} + \delta k_{23}k_{32} - \beta k_{23}k_{31} + \beta k_{21}k_{33} + \delta k_{33}k_{22} - \delta k_{22}k_{33});$
- (26) $e_3 \otimes e_2 \otimes e_3(-\beta k_{33}k_{13} + \beta k_{31}k_{33} - \delta k_{33}k_{23} + \delta k_{32}k_{33});$
- (27) $e_3 \otimes e_1 \otimes e_3(-k_{33}k_{13} + k_{31}k_{33});$
- Let $k_{11} = x, k_{22} = y, k_{33} = z, k_{12} = p, k_{21} = q, k_{13} = s, k_{31} = t, k_{23} = u, k_{32} = v.$
- It follows from (1)-(27) that
- (28) $-sx + xt = 0;$
- (29) $-\beta up + \beta qv - \delta uy + \delta yv = 0;$
- (30) $-vs + pz - \beta s^2 + \beta xz - \delta su + \delta zp = 0;$
- (31) $-\beta xz + \beta st - \delta zq + \delta ut - ut + qz = 0;$
- (32) $-zp + tv - \beta zx + \beta ts - \delta zp + \delta vs = 0;$
- (33) $-zp + sv - \beta st + \beta xz - \delta sv + \delta zp = 0;$
- (34) $-\beta zx + \beta tt - \delta zq + \delta vt - zq + tu = 0;$
- (35) $-\beta st + \beta xz - \delta tu + \delta qz - us + qz = 0;$
- (36) $-tp + xv - sp + vx = 0;$
- (37) $-ux + qt - ux + qs = 0;$
- (38) $-st + xz - s^2 + xz = 0;$
- (39) $-zx + tt - zx + st = 0;$
- (40) $-\beta vx + \beta pt - \delta vq + \delta yt - \beta ux + \beta qt - \delta uq + \delta yt = 0;$

- (41) $-\beta sp + \beta xv - \delta sy + \delta pv - \beta sq + \beta xu - \delta sy + \delta pu = 0;$
- (42) $-\beta vs + \beta pz - \delta vu + \delta yz - \beta su + \beta zq - \delta u^2 + \delta yz = 0;$
- (43) $-\beta zp + \beta tv - \delta zy + \delta vv - \beta zq + \beta tu - \delta zy + \delta vu = 0;$
- (44) $-vx + pt - \beta sx + \beta xt - \delta sq + \delta pt - sq + xu = 0;$
- (45) $-\beta pt + \beta vx - \delta ty + \delta qv - up + qv - \beta ux + \beta qs - \delta up + \delta ys = 0;$
- (46) $-\beta zp + \beta sv - \beta ut + \beta qz = 0;$
- (47) $-\beta zs + \beta tz - \delta zu + \delta vz = 0;$
- (48) $-zs + tz = 0;$

It is clear that r is a solution of CYBE iff (28)-(48) hold.

(I) It is trivial.

(II) Let $\beta = 0$ and $\delta \neq 0$. We only show that if r is a solution of CYBE, then r is one of two cases in part (II). If $z \neq 0$, we have that $u = v, s = t, yz = u^2, xz = s^2$ and $ux = sp$ by (47), (48), (42), (38) and (36), respectively. It follows from (30) and (33) that $pz = us$ and from (31) and (34) that $qz = us$. Thus $q = p$ and $xy = p^2$ since $xyz = xu^2 = spu = zp^2$. By Lemma 1.2, r is strongly symmetric.

If $z = 0$, we have that $u = -v$ and $s = -t$ by (43), (42), (38) and (39). It follows from (28), (29), (36), (40), (30), (44) and (45) that $sx = 0, yu = 0, xv = 0, ys = 0, (1 - \delta)us = 0, (1 + \delta)s(p + q) = 0$ and $(1 + \delta)u(p + q) = 0$, respectively. Consequently, r is the second case.

(III) Let $\beta \neq 0, \delta = 1$. We only show that if r is a solution of CYBE, then r is one case in part (III). If $z \neq 0$, we have that $t = s, u = v, q = p, xz = s^2, zp = su, sp = xu$ and $u^2 = yz$ by (48), (47), (46), (38), (32), (36) and (42), respectively. Since $xyz = xu^2 = spu = zp^2, xy = p^2$. Thus r is strongly symmetric by Lemma 1.2.

If $z = 0$, then $st = 0, t = 0$ and $s = 0$ by (31), (39) and (38), respectively. It follows from (42) and (43) that $u = -v$. By (36), (45) and (29), $xu = 0, u(p + q) = 0$ and $uy = 0$. Consequently r is the second case.

(IV) Let $\beta = \delta = 0$. We only show that if r is a solution of CYBE, then r is one case in part (IV). If $z \neq 0$, we have that $s = t, s^2 = xz, zp = vs$ and $zq = us$ by (48), (38), (32) and (34), respectively. Consequently r is the first case in part (IV). If $z = 0$, then $s = -t$ by (38) and (39). It follows from (30), (31), (28), (37), (36), (45) and (44) that $vs = 0, us = 0, xs = 0, xu = 0, xv = 0, up = qv$ and $s(p + q) = 0$. Consequently, r is the second case in part (IV). $\square$

Corollary 1.7 *Let L is a Lie algebra with $\dim L \leq 3$ and $r \in L \otimes L$. If r is strongly symmetric, then r is a solution of CYBE in L .*

Proof . If k is algebraically closed, then r is a solution of CYBE by Proposition 1.4, 1.6 and 1.3. If k is not algebraically closed, let P be algebraically closure of k . We can construct a Lie algebra $L_P = P \otimes L$ over P , as in [3, section 8]. Set $\Psi : L \longrightarrow L_P$ by sending x to $1 \otimes x$. It is clear that L_P is a Lie algebra over P and Ψ is homomorphic with $\ker \Psi = 0$ over k . Let

$$\bar{r} = (\Psi \otimes \Psi)(r).$$

Obviously, $\bar{r}$ is strongly symmetric. Therefore $\bar{r}$ is a solution of CYBE in L_P and so is r in L . $\square$

2 Coboundary Lie bialgebras

In this section, using the general solution, which are obtained in the section above, of CYBE in Lie algebra L with $\dim L \leq 3$, we give the the sufficient and necessary conditions which $(L, [\], \Delta_r, r)$ is a coboundary (or triangular) Lie bialgebra.

We now observe the connection between solutions of CYBE and triangular Lie bialgebra structures. It is clear that if $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra then r is a solution of CYBE. Conversely, if r is a solution of CYBE and r is skew symmetric with $r \in L \otimes L$, then $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra by [8, Proposition 1] .

Theorem 2.1 *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that $[e_1, e_2] = e_3, [e_2, e_3] = \alpha e_1, [e_3, e_1] = \beta e_2$, where $\alpha, \beta \in k$ and $\alpha\beta \neq 0$ or $\alpha = \beta = 0$. Let $p, u, s \in k$ and $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$. Then*

(I) *$(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff r is skew symmetric;*

(II) *$(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra iff*

$$\beta s^2 + \alpha u^2 + p^2 = 0,$$

Proof. (I) It is sufficient to show that

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_i) = 0$$

for $i = 1, 2, 3$. First, by computation, we have that

$$\begin{aligned} & (1 \otimes \Delta)\Delta(e_1) \\ &= \{(e_1 \otimes e_2 \otimes e_3)(\beta ps - \beta ps) + (e_3 \otimes e_1 \otimes e_2)(\beta ps) + (e_2 \otimes e_3 \otimes e_1)(-\beta sp)\} \\ &+ \{(e_1 \otimes e_3 \otimes e_2)(-\beta ps + \beta ps) + (e_3 \otimes e_2 \otimes e_1)(-\beta ps) + (e_2 \otimes e_1 \otimes e_3)(\beta sp)\} \\ &+ \{(e_1 \otimes e_1 \otimes e_2)(\alpha\beta su) + (e_1 \otimes e_2 \otimes e_1)(-\beta\alpha su) + (e_2 \otimes e_1 \otimes e_1)(0)\} \end{aligned}$$

$$\begin{aligned}
& + \{ (e_1 \otimes e_1 \otimes e_3)(-\alpha pu) + (e_1 \otimes e_3 \otimes e_1)(\alpha pu) + (e_3 \otimes e_1 \otimes e_1)(0) \} \\
& + \{ (e_2 \otimes e_1 \otimes e_2)(-\beta^2 s^2) + (e_1 \otimes e_2 \otimes e_2)(0) + (e_2 \otimes e_2 \otimes e_1)(\beta^2 s^2) \} \\
& + \{ (e_3 \otimes e_3 \otimes e_1)(p^2) + (e_3 \otimes e_1 \otimes e_3)(-p^2) + (e_1 \otimes e_3 \otimes e_3)(0) \}
\end{aligned}$$

Thus

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_1) = 0$$

Similarly, we have that

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_2) = 0$$

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_3) = 0$$

Thus $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra.

(II) By part (I), it is sufficient to show that r is a solution of CYBE iff

$$\beta s^2 + \alpha u^2 + p^2 = 0.$$

(a). Let $\alpha \neq 0$ and $\beta \neq 0$. Considering r is skew symmetric, by Proposition 1.4 (II), we have r is a solution of CYBE iff

$$\beta s^2 + \alpha u^2 + p^2 = 0.$$

(b). Let $\alpha = \beta = 0$. Considering r is skew symmetric, by Proposition 1.4 (III), we have r is a solution of CYBE iff

$$\beta s^2 + \alpha u^2 + p^2 = 0.$$

□

Let $(L, [\])$ be a familiar Lie algebra, namely, Euclidean 3-space under vector cross product. By Theorem 2.1, $(L, [\], \Delta_r, r)$ is not a triangular Lie bialgebra for any $0 \neq r \in L \otimes L$. In fact, [5, Example 2.14] already contains the essence of this observation.

Example 2.2 *Under Example 1.5, we have the following:*

- (i) $(sl(2), [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff r is skew symmetric;
- (ii) $(sl(2), [\], \Delta_r, r)$ is a triangular Lie bialgebra iff

$$-4s^2 + 4u^2 + p^2 = 0,$$

where $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$ and $p, s, u \in k$,

Theorem 2.3 *Let L be a Lie algebra with a basis $\{e_1, e_2, e_3\}$ such that*

$$[e_1 e_2] = 0, [e_1, e_3] = e_1 + \beta e_2, [e_2, e_3] = \delta e_2,$$

where $\delta, \beta \in k$. Let $p, s, u \in k$ and $r = p(e_1 \otimes e_2) - p(e_2 \otimes e_1) + s(e_1 \otimes e_3) - s(e_3 \otimes e_1) + u(e_2 \otimes e_3) - u(e_3 \otimes e_2)$

(I) $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra iff

$$(\delta + 1)((\delta - 1)u + \beta s)s = 0;$$

(II) If $\beta = 0$, then $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra iff

$$(1 - \delta)us = 0.$$

(III) If $\beta \neq 0$ and $\delta = 1$, then $(L, [\], \Delta_r, r)$ is a triangular Lie bialgebra iff

$$s = 0.$$

iff $(L, [\], \Delta_r, r)$ is a coboundary Lie bialgebra.

Proof . (I) We get by computation

$$\begin{aligned} & (1 \otimes \Delta)\Delta(e_1) \\ = & \{(e_1 \otimes e_1 \otimes e_2)(\beta\delta ss - \delta us) + (e_1 \otimes e_2 \otimes e_1)(-\beta\delta ss + \delta us) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\ & + \{(e_2 \otimes e_2 \otimes e_1)(-\beta us + uu + \beta^2 s^2 - \beta su) \\ & + (e_2 \otimes e_1 \otimes e_2)(\beta us - uu + \beta us - \beta^2 s^2) + (e_1 \otimes e_2 \otimes e_2)(0)\} \\ & (1 \otimes \Delta)\Delta(e_2) \\ = & \{(e_1 \otimes e_1 \otimes e_2)(\delta^2 ss) + (e_1 \otimes e_2 \otimes e_1)(-\delta^2 s^2) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\ & + \{(e_2 \otimes e_2 \otimes e_1)(\delta\beta ss - \delta su) + (e_2 \otimes e_1 \otimes e_2)(\delta su - \delta\beta ss) + (e_1 \otimes e_2 \otimes e_2)(0)\} \\ & (1 \otimes \Delta)\Delta(e_3) \\ = & \{(e_1 \otimes e_2 \otimes e_3)(\delta su + \beta ss) + (e_2 \otimes e_3 \otimes e_1)(-\delta us - \beta ss) \\ & + (e_3 \otimes e_1 \otimes e_2)(\delta\delta us + \beta\delta ss + \beta ss - su)\} \\ & + \{(e_1 \otimes e_3 \otimes e_2)(-\delta su - \beta ss) + (e_3 \otimes e_2 \otimes e_1)(-\beta\delta ss - \delta\delta us - \beta ss + su) \\ & + (e_2 \otimes e_1 \otimes e_3)(\delta us + \beta ss)\} \\ & + \{(e_1 \otimes e_1 \otimes e_2)(-\delta sp - \delta\delta ps + sp + \delta sp) \\ & + (e_1 \otimes e_2 \otimes e_1)(\delta ps + \delta\delta ps - \delta sp - sp) + (e_2 \otimes e_1 \otimes e_1)(0)\} \\ & + \{(e_1 \otimes e_1 \otimes e_3)(ss) + (e_1 \otimes e_3 \otimes e_1)(-ss) + (e_3 \otimes e_1 \otimes e_1)(0)\} \end{aligned}$$

$$\begin{aligned}
& + \{ (e_2 \otimes e_2 \otimes e_1)(-\beta ps + pu - \delta\beta sp - \delta\delta up - \delta\beta ps + \delta pu - \beta sp - \delta up) \\
& + (e_2 \otimes e_1 \otimes e_2)(\beta ps - pu + \delta\beta sp + \delta\delta up + \delta\beta ps - \delta pu + \beta sp + \delta up) \\
& - (e_1 \otimes e_2 \otimes e_2)(0) \} \\
& + \{ (e_2 \otimes e_2 \otimes e_3)(\delta\delta uu + \delta\beta us + \beta\delta us + \beta\beta ss) \\
& + (e_2 \otimes e_3 \otimes e_2)(-\delta\delta uu - \delta\beta us - \beta\delta us - \beta\beta ss) + (e_3 \otimes e_2 \otimes e_2)(0) \}
\end{aligned}$$

Consequently,

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_1) = 0$$

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_2) = 0$$

and

$$(1 + \xi + \xi^2)(1 \otimes \Delta)\Delta(e_3) = 0$$

iff

$$\delta^2 us + \delta\beta s^2 + \beta s^2 - us = 0.$$

This implies that $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff

$$\delta^2 us + \delta\beta s^2 + \beta s^2 - us = 0$$

(II) If $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra, then $(\delta^2 - 1)us = 0$ by part (I). If $(\delta + 1) \neq 0$, then $(1 - \delta)us = 0$. If $\delta + 1 = 0$, then $(1 - \delta)us = 0$ by Proposition 1.6 (II). Conversely, if $(1 - \delta)us = 0$, then we have that $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra by part (I). Since r is skew symmetric we have that r is a solution of CYBE by Proposition 1.6 (II) (IV). Thus $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra.

(III) Similarly, we can show that part (III) holds by part (I) and Proposition 1.6 (III).

□

Theorem 2.4 *If L is a Lie algebra with $\dim L = 2$ and $r \in L \otimes L$, then $(L, [\cdot, \cdot], \Delta_r, r)$ is a triangular Lie bialgebra iff $(L, [\cdot, \cdot], \Delta_r, r)$ is a coboundary Lie bialgebra iff r is skew symmetric.*

Proof. It is an immediate consequence of the main result of [5]. □

Acknowledgement I would like to show my gratitude to Professor Yonghua Xu and referee for their guidance and help.

References

- [1] *a.a. Belavin and V.G.Drinfel'd, solutions of the classical Yang-Baxter equations for simple Lie algebras, Functional anal. appl*, **16**, No. 3, 159-180.
- [2] *V.G.Drinfel'd, Quantum groups, in "Proceedings International Congress of Mathematicians, august 3-11, 1986, Berkeley, Ca" pp. 798-820, amer. Math. Soc., Providence, RI, 1987.*
- [3] *N.Jacobson, Lie algebras, Interscience publishers a division of John Wiley and Sons , New York, 1962.*
- [4] *S. Majid, Physics for algebraists: Non-commutative and non-cocommutative Hopf algebras by a bicrossproduct construction, J.Algebra* **130** (1990), 17-64.
- [5] *W. Michaelis, a class of infinite-dimensional Lie bialgebras containing the Virasoro algebra, advances in mathematics,* **107** (1994), 365-392.
- [6] *W. Michaelis, Lie coalgebras, Advances in mathematics,* **38** (1980), 1-54.
- [7] *W. Michaelis, The dual Poincare-Birkhoff-Witt Theorem, Advances in mathematics,* **57** (1985), 93-162.
- [8] *E.J.Taft, Witt and Virasoro algebras as Lie bialgebras, J.Pure Appl. bialgebra,* **87** (1993), 301-312.
- [9] *C.N.Yang and M.L.Ge, eds., Braid group, knot theory and statistical mechanics,* eds. C.N.Yang and M.L.Ge (World scientific , Singapore, 1989);